


Accession #:

Test Code: 4990

Patient Name:

Estrogen Ratios







µg/g Creatinine

µg/g Creatinine

L

144

60

Glucocorticoids

| Pregnanetriol | 193 |
|----------------------------------|------|
| Cortisone (E) | 79 |
| Cortisol (F) | 61 |
| Tetrahydrocortisone (THE) | 2098 |
| Allo-Tetrahydrocortisol (5α-THF) | 541 |
| Tetrahydrocortisol (THF) | 1118 |
| Adrenal Reserve (THE+5α-THF+THF) | 3757 |
| 11β-Hydroxyandrosterone | 620 |
| 11β-Hydroxyetiocholanolone | 266 |


Mineralocorticoids

Allo-Tetrahydrocorticosterone (5α-THB)

11-Dehydrotetrahydrocorticosterone (THA) 43

Tetrahydrocorticosterone (THB)

Reference Range



Reference Range



| Accession #: | Test Code: 4990 | | | |
|------------------------|-----------------------------------|--------------------|-----------------------------------|-------------|
| Patient Name: | | | | |
| Enzyme Activity Phenot | ype Assessment | | | |
| 5α-Reductase | | | 0.6 Referer | ace Range |
| Andro/Etio Ratio | 0.71 | | | |
| 5α-THF/THF Ratio | 0.48 | L | 0.6 | 1.6 |
| Elevated 5α-red | uctase activity is associated wit | ith Polycystic Ova | rian Syndrome (PCOS) and hirsutis | m in women, |

Benign Prostatic Hyperplasia (BPH) and premature baldness in men, and obesity and insulin resistance in both genders. Low 5α -reductase activity may result in reduced conversion of testosterone to DHT and undervirilization in males.

0.5

1.2

11β-HSD II (11β-hydroxysteroid dehydrogenase II)

Cortisol/Cortisone Ratio (118-HSD II) 0.77

11 β -HSD II is predominantly a renal enzyme. It inactivates cortisol in order to prevent competitive binding to mineralocorticoid receptors. Its activity can be measured by the ratio of cortisol/cortisone. An elevated ratio (toward right on the graph) indicates suppressed enzyme activity, and may be clinically related to stress, hypertension, high dose licorice, cortisol administration, or insulin resistance.

| Other Analytes | | | | |
|----------------------------------|-----------------|---------|----------|--|
| Melatonin | | | | |
| | μg/g Creatinine | Referen | ce Range | |
| | | 17.7 | 84.4 | |
| 6-Sulfatoxymelatonin (1st Mornir | ng) 62.2 | | | |
| Thyroid | | | | |
| | μg/g Creatinine | Referen | ce Range | |
| Free T3 | 339 | 266 | 938 | |
| The IS | | | | |
| Free T4 | 630 | 341 | 1524 | |
| | | | | |
| Organic Acids | | | | |
| | mg/g Creatinine | Referen | ce Range | |
| Kupurapia | 0.02 | 0.6 | 5 | |
| Kynurenic | 0.93 | | | |
| Xanthurenic | 0.33 | 0.2 | 1.5 | |
| | | | | |


| A | | ш. |
|------|-------|----|
| Acce | ssion | #: |


Test Code: 4990


Patient Name:

4-Point Cortisol and Cortisone

| Cortisol | µg/g | Range |
|-------------|--------|------------|
| 1st Morning | 23.1 | 4.4 - 26.9 |
| 2nd Morning | 14.6 L | 15.5 - 108 |
| Afternoon | 7.1 | 6.2 - 26.7 |
| Night | 3.1 | 1.8 - 17.7 |







----- Upper and lower limits of reference range.

| Cortisone | µg/g | Range |
|-------------|------|--------------|
| 1st Morning | 55.9 | 18.2 - 69.3 |
| 2nd Morning | 61.3 | 41 - 177.1 |
| Afternoon | 45.7 | 28.4 - 101.3 |
| Night | 23.3 | 10.3 - 58.8 |

| Creatinine | mg/dL | Range |
|-------------|-------|----------|
| 1st Morning | 108 | 35 - 270 |
| 2nd Morning | 117 | 35 - 270 |
| Afternoon | 110 | 35 - 270 |
| Night | 58 | 35 - 270 |

